



A. Fatemi, DDS, FICOI, FAGD

American Dental Association • Academy of General Dentistry

International Congress of Oral Implantologists • Fellow of International Congress of Oral Implantologists

Southern Maryland Dental Society • Fellow of Academy of General Dentistry

Consent for Oral Surgery

I hereby authorize the doctor to perform the following procedure(s):

I am aware that the procedure will involve the use of local anesthetic and/or nitrous oxide. I realize that there are risks involved in this operation and in the anesthesia or sedation process.

Certain complications and risks during surgery, regardless of the use of IV sedation or not, include but are not limited to:

- | | |
|--|--|
| 1. Bleeding and swelling | 7. Opening of the sinus (when removing upper teeth) |
| 2. Post-operative infection | 8. Jaw fracture |
| 3. Trismus (inability to open mouth wide) | 9. Numbness in lower lip, chin, or tongue following of lower teeth |
| 4. Post-operative nausea and vomiting | 10. Cardiac arrest |
| 5. Injury to adjacent teeth | 11. Temporomandibular joint soreness |
| 6. Decision to leave small root in jaw when removal would be detrimental | 12. Dry socket |
| | 13. Other: _____ |

I further understand that this is an elective procedure and other forms of treatment or no treatment at all are choices that I have. I am satisfied with the doctor's explanation of the operation, risks, and/or possible alternative treatment. I specifically consent to the procedure outlined above and I authorize the doctor to perform such other procedures as may be necessary in his/her judgment if such a need arises in the course of this treatment.

Initials



A. Fatemi, DDS, FICOI, FAGD

American Dental Association Academy of General Dentistry

International Congress of Oral Implantologists Fellow of International Congress of Oral Implantologists

Southern Maryland Dental Society Fellow of Academy of General Dentistry

Consent for Oral Surgery

Paresthesia

The removal of impacted teeth can cause several post-operative problems. After removal of your impaction, you will receive adequate information to help you deal with these discomforts. One possible complication that you should understand before your surgery is the phenomenon "Paresthesia". This is a temporary numbing of the lower lip or tongue similar to the feeling when you've had Novocain. The position of certain impactions may lead to this condition following their removal. There is no way to prevent Paresthesia. If it occurs, numbness or tingling will last an indefinite period of time. Recovery usually occurs within a few days or in some cases several months. On very rare occasions, this condition can be permanent.

The doctor has made no guarantee to me regarding the procedure outlined above.

I have read and understood this consent form.

Patient or authorized person to sign for patient

Date

Patient Name (print)

If not the patient, what is your relationship to the patient? _____

I have explained the condition, procedure, benefits, alternatives, and risks described on this form to the patient or representative.

Doctor Signature

Date